Milky Spore

Use Milky Spore to Kill Japanese Beetle Grubs

Japanese Beetle Larvae live in the soil feeding on tender grass roots.

1,000 Adult Beetle lay 50,000 Grubs each year.

Rid Your Lawn Of Unsightly Brown Spots



It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE CONTAINER TO FLUFF POWDER BEFORE USING.

Powder tends to settle during shipment.

FOR USE ON LAWNS AND GRASSY AREAS,
ORNAMENTAL GARDENS AND FLOWER BEDS
MULCHED REDS AND GROUND COVER REDS:

To control Japanese bestle grubs in Jawns and these other areas, apply one level measuring teaspoon of MILKY SPORE powder on top of grass or on the ground between plants, in spots every four feet along rows four feet apart. Use II 0 as to treat 2,500 sq.ft., I pound for 4,000 sq.ft., 2.5 lbs for I0,000 sq.ft., or ten pounds of MILKY SPORE powder per are of lawn.

MILKY SPORE can also be applied with the Milky Spore Dispenser Tube. Directions are included with the dispenser.

Treat your lawn or other areas specified on this label during any season of the year when the ground is not frozen and water lightly after application to soak spores into the soil.

Areas between treated spots will be inoculated naturally. Uniform, highly efficient control cannot be expected until the entire area has been inoculated. TEN YEAR GUARANTEE ON THIS PRODUCT WHEN APPLIED AS DIRECTED.









Magnified Spores



Grub infected with Milky Spore



Milky Spore
POWDER
JAPANESE BEETLE CONTROL



READY TO USE

NET WT. 2.5 LBS - TREATS 10,000 SQ. FT. BY SPOT APPLICATION
FPA REG. NO. 63191-1 - FPA EST. NO. 63191-VA-1

See other nanels (or product insert)

for precautionary statements, first aid, and directions for use

KEEP OUT OF REACH OF CHILDREN

Milky Spore

Use Milky Spore to Kill Japanese Beetle Grubs

PRECAUTIONARY STATEMENTS:

Hazards to Humans and Domestic Animals:

CAUTION: Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wath contaminated dothing before reuse.

ENVIRONMENTAL HAZARDS:

To protect the environment, do not allow pesticide to enter or run of first ost mort draits, draining effiches, gotters or surface waters. Applying this product in calm weather when heavy rain in our predicted for the not 24 hours will help to ensure that wind or rain does not blow or wish pasticide of the treatment area. Sweeping any product that had cold off the treatment area. Sweeping any product that had area of the laws or graters, with red prevent run off to water holds or of draines systems.

FIRST AID:

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treat-ment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment adules.

IF IN EYES:

Hold eye open and rinse slowly and gendy with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE:

Store in a cool, dry place away from direct sunlight. Keep container tightly closed to protect product from maisture.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill container.

IF EMPTY: Place in trash or offer for recycling.
 if available.

 IF PARTLY FILLED: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

HOTLINE NUMBER:

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information or other questions call 1-800-801-0061, Monday through Friday, 9 a.m. to 5 p.m., Eastern time. During other times, call your poison control center as 1-800-222-1222.

WARRANTY:

To the extent consistent with applicable law, St. Gabriel Organics makes no other express or implied warranties, either of merchantibility or fitness, for a particular use and shall not be liable for misuse or conditions beyond its control, such as handling or storage.

Milky Spore

Use Milky Spore to Kill Japanese Beetle Grubs IAPANESE BEETLE / GRUB LIFE CYCLE



1,000 Adult Beetle lay 50,000 Grubs each year.

FOR MORE INFORMATION, GO TO WWW.StGabrielOrganics.com or CALL 1-800-801-0061

FREE BROCHURE:

"Organizing a Community Campaign against the Japanese Beetle." MADE IN USA BY:

ST. GABRIEL ORGANICS 14044 Litchfield Dr., Orange, VA 22960 BATCH CODE

