

grubGONE!® G

SPREADER SETTINGS & APPLICATION GUIDE

Apply grubGONE! G at a rate of 3 lbs per 1,000 sq.ft.



Spreader Type	Speed	37 oz (2.3 lb) per 1,000 sq ft	55 oz (3.4 lb) per 1,000 sq ft	Swath	Pattern / Cone
Agrifab Rotary	3 mph	4	4.5	?	n/a
Anderson's Rotary Model 2000 SR	3 mph	K	L	11	8
Anderson's Model 2000 Rotary Spreader	3 mph	K	L	11	6
Anderson's SSD Drop Spreader	3 mph	4	4.75	n/a	n/a
Earthway Ev-N-Spred Rotary Model 2170	3 mph	12	14	8	n/a
Gandy Drop	3 mph	19	21	n/a	n/a
LESCO Rotary 020093	3 mph	E	F	7	Open Full
LESCO Rotary 705698	3 mph	12	14	7	Open Full
Prizelawn BF1 Pull behind rotary model	3 mph	H	I	10	n/a
Prizelawn Rotary Model CBR-III	3 mph	H	I	8	n/a
Scotts Precision Green drop	3 mph	4	4.75	n/a	n/a
Scotts R-8A	3 mph	J	K	11	6
Spyker Rotary Model 46-22	3 mph	3.6	3.9	?	n/a
Vicon (03) Series	4.5 mph	19	23	30	-
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm)	4.5 mph	4.5	5.25	34	III-C
Scotts® Recommended Spreaders		Setting		Width of Coverage	
Scotts® Rotary Spreaders (EdgeGuard® DLX, EdgeGuard® Mini)		3		5'	
Scotts® Elite Spreader		3 3/4		6'	
Scotts® Hand-Held Spreaders (manual models)		3		N/A	
Scotts® Wizz® Year-Round Spreader (battery-powered)		3 1/2		5'	

Prill size of the pellets is 150 SGN. If you do not see your specific spreader on this chart, refer to the spreader's product guide using the grubGONE! prill size as a reference. Please note: These are estimates only, actual spreader application rate may vary due to spreader wear, operator use and other environmental factors & variables present during application.

Spring & Summer Application

Apply grubGONE! G during springtime (shortly after the ground has thawed) to control overwintering grubs that are coming up near the surface of your lawn to feed on the roots. Applying grubGONE! G during mid-summer is the best time to control newly hatched grub larvae that are in their first weeks of development. Apply within 2 - 3 weeks of peak flight of adult beetles which is typically near the beginning of August.

Fall Application

grubGONE! G is also an excellent product for fall grub control if you miss the mid-summer application window. Beetles spend their summer eating and laying eggs and by August, small (1st instar) grubs appear below the surface of your turf, feeding on the roots. Eventually turf damage becomes noticeable in the fall as these grubs grow into larger 2nd & 3rd instar grubs. The good news is that grubGONE! G is a proven winner against these larger grubs. Meaning you can successfully treat for grubs well into the fall season; thus helping you lessen the impact grubs and beetles have on your yard next year.

If grub infestation is extensive, we recommend applying grubGONE! in 2 of the 3 application windows described above.

No Weeding Required!

Since grubGONE! G is non-toxic to bees, butterflies, ladybugs and other beneficials there is no need to remove dandelions or other flowering weeds from your turf before application.

Warmer Climates May Have Continued Grub Activity *

Warmer Climates May Have Continued Grub Activity *

Spring Through Late June Late June to Mid August August to Mid October *

* Apply while grub or beetle invader is feeding. Consult with your local nursery, cooperative extension service or local pest control professional for precise timing of application in your region.

Phyllom BioProducts Corporation | info@phyllom.com | PhyllomBioProducts.com | GrubGONE.com
 Manufactured by Phyllom BioProducts Corporation