

→ NO RESIDUE

← NO RINSE

Use Trifecta **CROP CONTROL**
for treating and preventing problem mites, aphids,
thrips, white flies, PM, botrytis and other pests
and fungal/mildew issues.

Trifecta.com.bz

ACTIVE INGREDIENTS: Thyme Oil 14%, Clove Oil 10%, Garlic Oil 9%,
Peppermint Oil 4%, Corn Oil 3%, Geraniol 3%, Citric Acid 2%, Rosemary Oil 2%

INERT INGREDIENTS: 53% Filtered Water, Soap, 2-Propanol, Vinegar
Total.....100%

Trifecta, LLC. - 516 West Central Avenue - South Williamsport, PA 17702 - 570-337-5125 - Trifecta.com.bz
Manufactured by Inventek Colloidal Cleaners, LLC.

THIS PRODUCT MAY BE USED ON CROPS INTENDED FOR HUMAN CONSUMPTION. THIS PRODUCT IS EXEMPT FROM
REGISTRATION WITH THE U.S. ENVIRONMENTAL PROTECTION AGENCY UNDER FIFRA SECTION 25(b) REGULATIONS.

CAUTION

STORE ONLY IN ORIGINAL CONTAINER IN A DRY PLACE
INACCESSIBLE TO CHILDREN AND PETS. DO NOT REUSE
EMPTY CONTAINERS. OFFER FOR RECYCLING IF AVAILABLE OR
DISCARD IN TRASH.

KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH
EYES, SKIN OR CLOTHING. WEAR EYE PROTECTION AND GLOVES
WHEN HANDLING. IF IN EYES, FLUSH WITH WATER FOR AT LEAST
15 MINUTES. IF ON SKIN, WASH WITH PLENTY OF SOAP AND
WATER. GET MEDICAL ATTENTION IF IRRITATION PERSISTS. DO
NOT APPLY THIS PRODUCT UNDILUTED.

NOT INTENDED FOR AQUATIC USE.



CROP CONTROL

- SAFE FOR ALL CROPS -



NET CONTENTS: 32 FL. OZ. (946 ML)

DIRECTIONS FOR USE → shake well before diluting.

FOR PREVENTATIVE MAINTENANCE:
Dilute 1/2 oz. of Crop Control
for every gallon of water. Apply
Crop Control using an atomizer,
misting system or a spraying
apparatus. Thoroughly wet the
plants until foliage is dripping
including the underside of the
leaves. Minor wilting of feeder
leaves may occur until the plant
is acclimated to Crop Control.
Preventative applications
should be applied 1 time per week
and on the same day each week.

FOR INFESTATIONS:

Dilute 2 oz. of Crop Control for
every gallon of water. Apply
Crop Control using an atomizer,
misting system or a spraying
apparatus. Thoroughly wet the
plants until foliage is dripping
including the underside of the
leaves. Minor wilting of feeder
leaves may occur until the plant
is acclimated to Crop Control.
Crop Control should be applied
every other day up to 3 times
per week until infestation is
controlled. Apply Crop Control
at the end of a light cycle or
just before dusk to prevent
foliar damage.

