ARBER Bio Funigicide

For Use on Labeled:

Fruits, Vegetables, Nuts, Berries, Houseplants, Flowers, Foliage, Ornamental Trees, Turf, and Shrubs.

How To Use:

1 fluid oz (4 capfuls) of Arber Bio Fungicide per gallon of water. Spray plants to complete wetness, covering both top and bottom leaf surfaces to ensure complete coverage.

Hose End Sprayer:

Set dial on sprayer to deliver a rate of loz (2 tbsp) per gallon of water

Refer to booklet for complete directions for use

Active Ingredient:

Bacillus amyloliquefaciens strain F727*cells and spent fermentation media: 96.4% Other Ingredients: 3.6% Total:

*Contains a minimum of 1 X 10° cfu/mL of product

Distributed by: Evergreen Way Inc.

5999 Custer Rd. Ste 110-148 Frisco, TX 75035



EPA Rea. No: 84059 - 28 - 97021

FPA Est No: Printed On Container



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS: To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to apply this product in a manner inconsistent with its labeling.

ARBER Bio Fungicide is a broad-spectrum fungicide used for the control or suppression of a broad range of foliar diseases. ARBER Bio Fungicide may be used on vegetable crops, roses, fruits, nuts, flowers, foliage, houseplants, trees and shrubs located in residential landscapes. ARBER Bio Fungicide can be applied up to and on the day of harvest on all fruits and vegetables.

WHEN TO USE

For best results, apply ARBER Bio Fungicide prior to disease development or at the first sign of diseases and continue applying on a 7- to 10-day schedule until disease pressure subsides.

BEFORE YOU USE

Read and follow these directions when using. Do not allow spray to drift from application site. Use only with pressurized hand-held sprayers, spray trigger bottles, and hose-end sprayers. Do not allow spray mixture to stand overnight or for prolonged periods.

ARBER Bio Fungicide can be applied in commonly used pressurized hand-held sprayers, spray trigger bottles, and hose-end sprayers.

HOW TO USE FOR HAND-HELD SPRAYERS AND SPRAY TRIGGER BOTTLES

- Shake well before use.
- Fill sprayer or bottle with appropriate amount of water and concentrate.
- · Mix the spray solution thoroughly.
- Keep the spray solution agitated during application.

HOW TO USE FOR HOSE-END SPRAYERS

- Shake well before use.
- Follow hose-end sprayer instructions to determine how to fill, set dial clean and disconnect from hose
- Set dial on sprayer to deliver rate as directed below.

. .

HOW MUCH TO USE FOR ALL APPLICATIONS:

1 fluid ounce (2 TBSP) of ARBER Bio Fungicide per gallon of water.

Spray plants to complete wetness, covering both top and bottom leaf surfaces to ensure complete coverage.

Some pesticides can cause phytotoxic effects ranging from slight burning or browning of leaves to distorted leaves, fruit, flowers or stems. Damage symptoms may vary with the type of plant that has been treated. It is impossible to test all plant species for phytotoxicity. To assure that the plants to be treated are not sensitive

to the treatment, apply a small amount of the product to a few leaves or the above ground portion of the plant and check back in 2 - 4 days for signs of phytotoxicity. Use product according to label directions.

DISEASES CONTROLLED OR SUPPRESSED ON VEGETABLES. FRUITS, NUTS, ORNAMENTAL PLANTS*, HEMP, TOBACCO*, TREES, SHRUBS, FLOWERS INCLUDING HOPS, FOLIAGE, GREEN. AND TROPICAL PLANTS*

Anthracnose (Colletotrichum spp.)*

Bacteria (Erwinia spp., Pseudomonas spp., Xanthomonas spp.)

Bacterial Leaf Blight (Xanthomonas campestris)

Bacterial Speck (Pseudomonas syringae pv. Tomato)

Bacterial Spot (Xanthomonas spp.) Bean Rust (Uromyces appendiculatus)*

Black Mold (Alternaria alternata)*

Black Rot/Black Crown Rot (Alternaria spp.)*

Black Spot of Rose (Diplocarpon rosea)*

Botrytis (Botrytis spp.)*

Botrytis Leaf Blight (Botrytis squamosa)*

Botrytis Neck Rot (Botrytis spp.)*

Downy Mildew (Bremia lactucae, Peronospora spp., and Plasmopara viticola) - suppression

Early Blight (Alternaria solani)

Fire Blight (Erwinia amylovora) - suppression*

Grav Mold (Botrytis cinerea)*

Greasy Spot (Mycosphaerella citri)*

Late Blight (Phytophthora infestans) - suppression

Leaf Spots (Alternaria spp., Cercospora spp. Septoria spp.)* Onion Downy Mildew (Peronospora destructor)

Onion Purple Blotch (Alternaria porri)

Pin Rot (Alternaria/Xanthomonas complex)

Powdery Mildew (Uncinula necator, Erysiphe spp., Sphaerotheca spp., Oidiopsis taurica, Leveillula taurica, Podosphaera leucotricha, Oidium spp., Podosphaera spp.)
Rust (Puccinia spp.)*
Soch (Vonturia spp.)

Scab (Venturia spp.) - suppression* Sclerotinia Head and Leaf Drop (Sclerotinia spp.) - suppression Sour Rot (Alternaria tenuis, Aspergillus spp., Botrytis cinerea, Cladosporium herbarum, Penicillium spp., Rhizopus arrhizus)*

Cladosporium herbarum, Penicillium spp., Rhizopus arrhizus)* Target Spot (Corynespora cassiicola) Walnut Bliaht (Xanthomonas campestris)

White Mold (Sclerotinia sclerotiorum and S. minor)
*Not labeled for this use in California

STORAGE AND DISPOSAL Do not contaminate, water food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place, Avoid freezing

Pesticide Disposal and Container Handling: Nonrefilable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or (800) 858-7378 (National Pesticide InformationCenter) for disposal instructions. Never place unused product down any indoor or outdoor drain

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vorniting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Plinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible Call a poison control center or doctor for treatment advice.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For non-emergency information on product usage for example, call (1-877-664-4476), Monday through Friday, 9a.m. to 5 p.m Pacific Time. For medical emergencies call your poison control center at 1-800-222-1222

WARRANTY

To the extent consistent with applicable law, the seller makes no warranty, expressed or implied, of merchantability, fitness or otherwise concerning use of this product. To the extent consistent with applicable law, the user assumes all risks of use, storage or handling that are not in accordance with the accompanying directions.

US Patent No. 4.863.734 & No. 5.989.429

See attached booklet for complete direction for use and precautionary statements

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place. Avoid freezing

Pesticide Disposal and Container Handling: Nonrefillable container Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or (800) 858-7378 (National Pesticide InformationCenter) for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIDCT VID

FIRST AID	
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air if person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advise.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For non-emergenevinformation on product usage for example call (1-877-664-4476) Monday through Friday 9a m to 5 nm Pacific Time For medical emergencies call your poison control center at 1-800-222-1222

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eve irritation. Avoid contact with eves. skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse

ENVIRONMENTAL HAZARDS: To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

EPA Reg. No.: 84059-28 - 97021 FPA Est No. Printed on Container Lot No. printed on container

Distributed by: Evergreen Way Inc. 5999 Custer Rd. Ste 110-148 Frisco, Tx 75035