



PRESTOP®
WG



Agricultural, Turf and Ornamental Uses
For Use in Organic Production

Active Ingredient: *Gliricladium catenulatum* Strain J1446*93%
Other Ingredients:7%
Total:100%

* Contains a minimum of 1×10^9 cfu/g.

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID

If swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION:
Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Mixer/loaders and applicators must wear a NIOSH approved particulate respirator with any N, R, or P filter with NIOSH approval number prefix TC-84A; or a NIOSH approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS: When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides

[40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS: For terrestrial uses - Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Manufactured for:
Danstar Ferment AG
Poststrasse 30
CH-6300 ZUG
Switzerland

Marketing Company:
Lallemand Specialties Inc.
6120 West Douglas Avenue
Milwaukee, WI 53218
United States
www.lallemandplantcare.com



EPA Reg. No.: 64137-13

EPA Est. No.:

NET WEIGHT: 2.2 lbs (1 kg)



PRESTOP®
WG

Agricultural, Turf and Ornamental Uses
For Use in Organic Production

Active Ingredient: *Gliricladium catenulatum* Strain J1446*93%
Other Ingredients:7%
Total:100%

* Contains a minimum of 1×10^9 cfu/g.

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID

If swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.

Manufactured for:
Danstar Ferment AG
Poststrasse 30
CH-6300 ZUG
Switzerland

Marketing Company:
Lallemand Specialties Inc.
6120 West Douglas Avenue
Milwaukee, WI 53218
United States
www.lallemandplantcare.com



EPA Reg. No.: 64137-13

EPA Est. No.:

Lot No:

Use Before:

NET WEIGHT: 2.2 lbs (1 kg)

85A14-83-B6/17

Peel back for instructions. PRESS TO RESEAL.



Agricultural, Turf and Ornamental Uses

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION: Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Mixer/loaders and applicators must wear a NIOSH approved particulate respirator with any N, R, or P filter with NIOSH approval number prefix TC-84A; or a NIOSH approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS: When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS: For terrestrial uses - Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

EXCEPTION: If the product is soil incorporated or soil injected, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

USE INSTRUCTIONS

Apply Prestop (WG) by spraying or drenching the growth substrate, as a foliar spray, or incorporated into the growth substrate as an aqueous suspension.

Tank Mixes: Do not tank mix Prestop (WG) with any pesticides or concentrated fertilizers.

Integrated Pest Control Programs: Prestop (WG) can be used in integrated pest control programs. Always check the compatibility of Prestop (WG) with other Biological Fungicides or chemical pesticides with the local distributor. Do not apply chemical pesticides within 1 to 4 days of Prestop (WG) application except for propamocarb hydrochloride, tridemefon, deltamethrin, malathion, metalaxyl, pirimicarb and pyrethrins; these may be applied on the same day as Prestop (WG).

PRODUCT INFORMATION

Prestop (WG) controls seed-borne and soil-borne plant diseases such as damping-off, root and stem rot, and wilt caused by *Alternaria*, *Bipolaris*, *Botrytis*, *Cladosporium*, *Colletotrichum*, *Fusarium*, *Mycosphaerella*, *Penicillium*, *Phytophthora*, *Plasmiodiophora*, *Plicaria*, *Pyrenochaeta*, *Pythium*, *Rhizoctonia*, *Sclerotinia* and *Verticillium*, certain storage diseases caused by *Helminthosporium* and *Rhizoctonia*, and foliar diseases like grapevine trunk diseases (several pathogens) as well as diseases caused by *Alternaria*, *Anthraco**se* (*Colletotrichum*) *Bipolaris*, *Botrytis*, *Cladosporium*, *Didymella*, *Fusarium*, *Monilinia*, *Mycosphaerella*, *Penicillium* and *Sclerotinia* on greenhouse or field grown vegetables, herbs, ornamentals, berries, cereals, legumes, pome and stone fruits, nut trees, tree and forest seedlings and turf. Prestop (WG) also suppresses powdery mildew symptoms on these greenhouse or field grown crops.

Use Prestop (WG) on following agricultural grown crops in greenhouses or outdoors:

VEGETABLES

Fruiting, Leafy, Root Crop and Seed/Pod Vegetables including, but not limited to:

- | | | |
|--------------------|-----------------------|----------------|
| • Artichokes | • Cucumbers | • Parsnips |
| • Asparagus | • Eggplant | • Peas |
| • Beans | • Endive | • Peppers |
| • Beets | • Garlic | • Potatoes |
| • Broccoli | • Horseradish | • Pumpkins |
| • Brussels Sprouts | • Kale | • Radishes |
| • Cabbage | • Kohlrabi | • Rutabagas |
| • Carrots | • Leeks | • Spinach |
| • Cauliflower | • Lentils | • Squash |
| • Celery | • Lettuce | • Strawberries |
| • Chicory | • Melons | • Tomatoes |
| • Chinese Cabbage | • New Zealand Spinach | • Turnips |
| • Corn | • Onions | • Watermelons |

BERRIES including, but not limited to:

- | | |
|----------------|----------------|
| • Blueberries | • Grapevine |
| • Cane berries | • Strawberries |

POME AND STONE FRUITS including, but not limited to:

- | | |
|------------|-----------|
| • Apples | • Pears |
| • Apricots | • Peaches |
| • Cherries | • Plums |
| • Citrus | |

HERBS AND SPICES including but not limited to:

- | | | |
|-----------|------------|--------------|
| • Aniseed | • Ginseng | • Sage |
| • Basil | • Lavender | • Savory |
| • Caraway | • Marjoram | • Thyme |
| • Chives | • Oregano | • Watercress |
| • Dill | • Parsley | |
| • Fennel | • Rosemary | |

ORNAMENTALS

Flowers, Foliage and Potted, Bedding and Seedling Plants including, but not limited to:

- | | | |
|------------------|-----------------|----------------|
| • Achillea | • Delphinium | • Monarda |
| • African Violet | • Dianthus | • Nigella |
| • Ageratum | • Dieffenbachia | • Pansy |
| • Alyssum | • Freesia | • Pelargonium |
| • Amaryllis | • Fuchsia | • Petunias |
| • Anemone | • Gaillardia | • Phlox |
| • Anthurium | • Gazania | • Poinsettia |
| • Aster | • Geraniums | • Poppy |
| • Azalea | • Gerbera | • Primrose |
| • Begonias | • Gladiolus | • Rhododendron |
| • Calceolaria | • Rose | • Rose |
| • Campanula | • Gypsophila | • Rudbeckia |
| • Carnation | • Heder | • Salvia |
| • Centaurea | • Hyacinth | • Senecio |
| • Cerastium | • Impatiens | • Sinningia |
| • Chrysanthemum | • Iris | • Statice |
| • Cineraria | • Kalanchoe | • Sweet Pea |
| • Coleus | • Liatris | • Tulips |
| • Cyclamen | • Lilies | • Verbena |
| • Daffodils | • Lobelia | • Vinca |
| • Dahlia | • Marigolds | • Zinnia |
| • Daisy | • Matthiola | |

LEGUMES, CEREALS AND OIL SEED CROPS including, but not limited to:

- | | | |
|-----------|-----------|------------|
| • Alfalfa | • Canola | • Soybeans |
| • Barley | • Peanuts | • Wheat |

NUT TREES including, but not limited to:

- | | | |
|-----------|------------|-----------|
| • Almonds | • Chestnut | • Pecans |
| • Cashew | • Coconut | • Walnuts |

TOBACCO

TREES AND SHRUB SEEDLINGS including, but not limited to:

- | | | |
|----------------|--------------|---------------|
| • Birch | • Pine | • Woody |
| • Forest Trees | • Shrubs | • Ornamentals |
| • Fruit Trees | • Spruce | |
| • Juniper | • Vine Crops | |

TURF including, but not limited to:

- | | |
|--------------------------------|-------------|
| • Turf, grown for seed and sod | • Sod farms |
| • Turf, greens | • Lawns |

PREPARING AN AQUEOUS SUSPENSION OF Prestop (WG)

To prepare an aqueous suspension of Prestop (WG), mix product in a small volume of water such as 1/2 gallon (0.35 oz. of Prestop (WG) per 1/2 gallon of water). For better results, let the suspension stand 30 minutes. Agitate

sufficiently to evenly disperse product before diluting to final volume. Apply the aqueous suspension as a drench or spray or via drip irrigation or incorporation in potting media. Use the aqueous suspension within 1 day of preparation.

Examples of mixing ratios of Prestop (WG) with water are as follows:

Concentration of PRESTOP (WG) Suspension	Ounces (oz) of PRESTOP (WG)	Gallons of Water	Grams (g) of PRESTOP (WG)	Liters (L) of Water
0.01%	0.33	25	10	100
0.05%		5		20
0.1%		2.5		10
0.25%		1		4

INCORPORATION IN POTTING MEDIA

Incorporation of Prestop (WG) in potting media for vegetables, herbs, ornamentals and tree or forest seedlings grown in greenhouses or indoors. Mix Prestop (WG) in potting mixture at sowing, potting or transplanting as an aqueous suspension at the rate of 0.54 – 1.35 oz. of Prestop (WG) per cubic yard of growth substrate (i.e. 0.02 – 0.05 grams of Prestop (WG) per 1 liter of growth substrate).

When incorporating Prestop (WG) as an aqueous suspension, mix 0.33 oz. of product per 5 gallon of water (0.05% suspension). Use 8.1 – 20.4 gallons of 0.05% Prestop (WG) suspension per cubic yard of growth substrate (i.e. 0.02 – 0.05 grams of Prestop (WG) per liter of growth substrate).

Use the higher rate for higher disease pressure. It is important to evenly incorporate Prestop (WG) to assure best performance. Store potting media containing Prestop (WG) below 77°F (25°C) and use within 3 weeks after preparation. Incorporation of Prestop (WG) may be followed with a drench application within 3 – 6 weeks or sooner depending on disease pressure and new plant growth.

HYDROPONIC APPLICATIONS

Apply Prestop (WG) in hydroponic NFT, ebb-and-flow and rockwool applications according to following rates:

- NFT and ebb-and-flow application: 0.09 oz Prestop (WG) per 1000 plants, 0.002 oz/100 sq. ft. (i.e. 2.5 grams Prestop (WG) per 1000 plants; 0.05 kg/ha).
- Rockwool: 0.7 - 0.9 oz Prestop (WG) per 1000 plants, 0.013-0.016 oz/100 sq. ft. (i.e. 20-25 g/1000 plants, 0.4-0.5 kg/ha).
- Repeated hydroponic applications may be done within 3 – 6 weeks or sooner depending on disease pressure and new plant growth.

SOIL DRENCH APPLICATIONS

Use a 0.01 – 0.1% Prestop (WG) suspension (0.033-0.33 oz. of PRESTOP (WG) per 2.5 gallons of water) for soil drench applications. Refer to the above table for mixing ratios

Seedling Trays or Beds less than 4 inches deep (Recommended Rate per Area): Drench seedling trays or beds with Prestop (WG) at the rate of 0.35 oz. of Prestop (WG)/100 sq. ft. (1g/m2). Use a 0.05% Prestop (WG) suspension at 5 - 10 gal/100 sq. ft. Use the higher rate for higher crop density or higher disease pressure. For example, use 0.35 oz. of Prestop (WG)/5 gallons of water/50 sq. ft.

Pots, Plugs or Rockwool Cubes (Recommended Rate per 100 Plants): Drench crops grown in pots or rockwool cubes with Prestop (WG) using e.g.

a 0.05 % suspension. Dose rate depends upon the root plug/ball size and disease pressure. Calculate the amount of Prestop (WG) per plant based on the following recommendation: 0.54 – 1.35 oz. of Prestop (WG) per cubic yard of growth substrate (i.e. 0.02 – 0.05 grams of Prestop (WG) per liter of growth substrate). Use the higher rate for bigger root volume and higher disease pressure. Refer to the following table for application rates on different pot or root plug sizes:

Pot Size (Root Volume)	Ounces of Prestop (WG) per 1000 plants (oz per 1000 plants)	0.05% Prestop (WG) Suspension (fl. oz. per 1000 plants)
0.5 fl. oz. (15 ml)	0.01 – 0.03	20 – 51
2.0 fl. oz. (60 ml)	0.04 – 0.11	81 – 203
4.0 fl. oz. (120 ml)	0.09 – 0.21	162 – 406
10 fl. oz. (295 ml)	0.21 – 0.52	399 – 997
17 fl. oz. (500 ml)	0.35 – 0.88	676 – 1690
34 fl. oz. (1 L)	0.71 – 1.76	1353 – 3381
4 pints (1.9 l)	1.35 – 3.34	2571 - 6424

Timing of Treatments: Apply the first drench treatment at sowing. Repeat treatment at transplanting or potting. Repeated drench applications may be done within 3 – 6 weeks or sooner depending on disease pressure and new plant growth.

FOLIAR SPRAY

Spray or spread an aqueous suspension of Prestop (WG) on plant stems or foliar parts for control of *Alternaria*, *Anthraco**se* (*Colletotrichum*) *Bipolaris*, *Botrytis*, *Cladosporium*, *Didymella*, *Fusarium*, *Monilinia*, *Mycosphaerella*, *Penicillium*, *Sclerotinia* or grapevine trunk diseases (several pathogens). Use a water suspension of 0.33 oz. Prestop (WG) per 5 gallons water (0.05% suspension). Treat plants to the point of run-off using 0.05% Prestop (WG) suspension.

Use Prestop (WG) at 0.01-0.016 oz. / 100 sq. ft. for fogging applications of greenhouse grown crops such as vegetables, berries and ornamentals together with a minimum water volume of 0.005 gallon per 100 sq. ft. (equivalent to 0.3-0.5 kg Prestop (WG) / ha and min. water volume of 20 L/ha).

Repeat foliar spray / fogging every 3 – 6 weeks if necessary, or sooner depending on disease pressure and new plant growth.

Surfactants are not generally needed. A non-ionic surfactant may be used in combination with Prestop (WG) for crops with waxy / hard-to-wet leaf surfaces.

TREATMENT OF CUTTINGS, BULBS OR TUBERS

Dip rooted cuttings in a suspension of 0.1 oz. Prestop (WG) in one gallon of water (0.075% suspension) or spray with the suspension after planting. Dip or spray bulbs and tubers with a 0.075% Prestop (WG) suspension before planting or storage.

PREPARING AN AQUEOUS SUSPENSION OF Prestop (WG)

To prepare an aqueous suspension of Prestop (WG), mix product in a small

volume of water such as 1/2 gallon (0.35 oz. of Prestop (WG) per 1/2 gallon of water). For better results, let the suspension stand 30 minutes. Agitate sufficiently to evenly disperse product before diluting to final volume. Apply the aqueous suspension as a drench or spray or via drip irrigation or incorporation in potting media. Use the aqueous suspension within 1 day of preparation.

Examples of mixing ratios of Prestop (WG) with water are as follows:

Concentration of PRESTOP (WG) Suspension	Ounces (oz) of PRESTOP (WG)	Gallons of Water	Grams (g) of PRESTOP (WG)	Liters (L) of Water
0.01%	0.33	25	10	100
0.05%		5		20
0.1%		2.5		10
0.25%		1		4

SOIL DRENCH APPLICATIONS

Use a 0.01 – 0.1% Prestop (WG) suspension (0.033 - 0.33 oz. of PRESTOP (WG) per 2.5 gallons of water) for soil drench applications. Refer to the above table for mixing ratios.

FOLIAR SPRAY

Spray or spread an aqueous suspension of Prestop (WG) on plant stems or foliar parts for control of *Alternaria*, *Anthraco**se* (*Colletotrichum*) *Bipolaris*, *Botrytis*, *Cladosporium*, *Didymella*, *Fusarium*, *Monilinia*, *Mycosphaerella*, *Penicillium*, *Sclerotinia* or grapevine trunk diseases (several pathogens). Use a water suspension of 0.33 oz. Prestop (WG) per 5 gallons water (0.05% suspension). Treat plants to the point of run-off using 0.05% Prestop (WG) suspension.

Use Prestop (WG) at 0.01-0.016 oz. / 100 sq. ft. for fogging applications of greenhouse grown crops such as vegetables, berries and ornamentals together with a minimum water volume of 0.005 gallon per 100 sq. ft. (equivalent to 0.3-0.5 kg Prestop (WG) / ha and min. water volume of 20 L/ha).

Repeat foliar spray / fogging every 3 – 6 weeks if necessary, or sooner depending on disease pressure and new plant growth.

Surfactants are not generally needed. A non-ionic surfactant may be used in combination with Prestop (WG) for crops with waxy / hard-to-wet leaf surfaces.

CHEMIGATION USE DIRECTIONS

For Vegetable and Ornamental Crops:

- On an area basis, use Prestop (WG) at 0.08–0.16 oz. /100 sq. ft. (2.5 – 5 g/10 sq. meters).
- On an individual plant basis, use Prestop (WG) at 0.22 oz. per 250 plants.
- Use higher rate and more frequent application for large plants and pots, and for high disease pressure.
- Apply in sufficient amount of water to move into root zone. Repeat every 3 – 6 weeks as needed for disease control, or sooner depending on disease pressure and new plant growth.

General Requirements:

1. Apply this product only through drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

3. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Requirements for Chemigation Systems Connected to Public Water Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

4. The pesticide injection pipeline must contain a functional, normally closed, solenoidoperated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Drip (Trickle) Chemigation:

1. The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

3. The pesticide injection pipeline must also contain a functional, normally closed, solenoidoperated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being filled with a system interlock.

Application Instructions for All Types of Chemigation:

1. Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residues may cause product to lose effectiveness or strength.

2. Determine the treatment rates as indicated in the directions for use and make proper dilutions.

3. Prepare a solution in the chemical tank by filling the tank with the required water and then adding product as required. Utilize agitation to keep solution in suspension.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Prestop (WG) consists of living microbes that are packed in unit packages. Store unopened packages up to 12 months in a cool (below 41°F or 5°C), dry place and up to 2 weeks at room temperature (maximum 77°F or 25°C). Immediately use the entire package's content after opening.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Non refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration. Do not burn, unless allowed by state and local ordinances. (For instances where state and local ordinances do allow burning): If burned, stay out of smoke.

NOTICE TO USER

Lallemand Specialties Inc. warrants only that this product conforms to the product description on this label and is reasonably fit for the purposes set forth in the Directions for Use when used in accordance with them. However, ineffectiveness or other unintended consequences may result because of such factors as the use, storage or handling of the product contrary to the label instructions, all of which are beyond the control of Lallemand Specialties Inc. To the extent permitted by state law, Lallemand Specialties Inc. shall not be liable for indirect or consequential damages resulting from the use, storage or handling of this product. LALLEMAND SPECIALTIES INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.