

Mix Rate: The mix rate for centrate is 1.0 fl. oz. (2 tbs.) or 29.6 mL per gallon of water. For use only as a foliar spray!

Dilute and evenly apply SMITE to the top and underside of the leaves until all foliage is saturated. Start from the lowest part of the canopy and move to the top until all foliage is evenly coated.

- · Effective against all spider mites and on any crop, lawn, ornamental. or turf
- Tested on Tetranychus Urticae, which attacks beans, corn, peppers, potatoes, strawberries, tomatoes, and ornamentals such as roses.
- · Effective for spider mite control, broadly recommended, and used in various fields, such as forestry, horticulture, lawn care, and organic farming.

56360"00499



### **ACTIVE INGREDIENTS:**

Geraniol	7.680%
Peppermint oil	0.256%
Cottonseed oil	0.026%
Rosemary oil	0.003%
OTHER INGREDIENTS:*	
TOTAL:	100.000%

\*Water, Canola oil, Polyoxyethylene lauryl ether

## **FIRST AID**

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor. . Do not give anything by mouth to an unconscious person. IF INHALED . Move person to fresh air. . If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth to mouth if possible. . Call a poison control center or doctor for further treatment advice. IF IN EYES . Hold eve open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eve. • Call a poison control center or doctor for treatment advice. IF ON SKIN OR CLOTHING . Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment. HOT LINE NUMBER: Have the product container or label with vou when calling a poison control center or doctor, or going for treatment. For emergency information call the National Pesticides Information Center at 1-800-858-7378. 6:30 AM to 4:30 PM Pacific time (PT), seven days a week. During other times, call the poison control center 1-800-222-1222

### **DIRECTIONS FOR USE**

Phytotoxicity: Plants may be sensitive to this product. Always test a small area on a plant before making a broad application.

Preparation: concentrate is 1.0 fl. oz. (2 tbs.) or 29.6 mL per gallon of water. See chart below. Add concentrate

to water. Shake and mix vigorously.

Application: For optimum results, shake and mix vigorously, apply during the early morning or after peak sunlight and before sundown. Do not use in direct sunlight or intense artificial lighting. Spray directly on infested foliage. Apply until all foliage is evenly covered. Apply to infested area every 5-7 days until infestation is controlled

every 10-14 days to prevent insect infestation.

Usage	Mix Rate	Dilution	Foliar Spray	Timing & Frequency
All crops, lawns, or ornamentals	1.0 fl. oz. (29.6 mL) / 1 gal.	1:128	3.0 fl. oz. (88.7 mL) / 1,000 sq. ft.	Apply every 5-7 days during spider mite infestation or every 10-14 days as a preventative measure.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

Pesticide Storage: Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

# KEEP OUT OF REACH OF CHILDREN See Back Panel for Precautionary Statements

NET CONTENTS: 32 fl. oz. (946 mL)

Supreme Growers LLC 57 Cool Springs Road Signal Mountain, TN 37377 (423) 803-2500

is extremely effective and kills

naturally. It contains oils produced by

cold-pressed extraction of naturally

grown Geranium, Peppermint, Cotton-

seed, and Rosemary. Under warm con-

ditions, SMITE will naturally biodegrade

Spider mites are known to feed on sev-

eral hundred species of plants. A sin-

gle mature female can spawn a million

mites in one month. Their accelerated

reproductive rates allow spider mites

to adapt quickly and develop resistance

blocking the breathing holes, resulting

reproductive disruption among female

spider mites not initially contacted by

the spray. This leads to infertility, eggs

that do not hatch or mites that do not

leaving minimal residue.

to normal pesticides.

in death by suffocation.

develop to an adult stage.

SUPREMEGROWERS® SUPREME GROWERS LLC. ©Copyright 2010-2018

### ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

NON-TOXIC • SUPREME QUALITY • ENVIRONMENTALLY SAFE •

Smite 32OZ Bottle Label 7176CR1.indd 1 2/21/18 1:57 PM