

BIOMIN[®] BORON

1-0-0

Mineral for Soil and Foliar Application



Product Information

BIOMIN BORON contains boron for soil and/or foliar application. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct boron deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizers.

GUARANTEED ANALYSIS

Total Nitrogen (N).....1.0%
1.0% Water Soluble Nitrogen
Boron (B)3.0%

Derived from sodium tetraborate and hydrolyzed vegetable protein.

KEEP OUT OF REACH OF CHILDREN

CAUTION - This product contains Boron. Do not apply to Boron sensitive crops. Use of the product on any crop other than those recommended may result in serious injury to the crop(s).

SHAKE WELL BEFORE USE

PREVENTS AND CORRECTS BORON DEFICIENCIES

NET CONTENTS: 2.5 Gallons (9.45 liters)

Weight per Gallon: 9.5 lb (4.3 kg) @ 68°F

U.S. Pat.# 5,504,055

Lot #:



Directions and General Recommendations

SOIL APPLICATION

BIOMIN BORON may be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1 quart-1 gallon per acre (2.5-10 liters per hectare) depending on crop, soil conditions and application methods. For maintenance, 1 quart per acre (2.5 litres per hectare) is recommended. **BIOMIN BORON** should be diluted with enough water to cover entire acre.

FOLIAR APPLICATION

FIELD CROPS: Apply 1-4 pints per acre (1.25-5 liters per hectare) of **BIOMIN BORON** when crops are 6-12 inches (15-30 cm) in height or when deficiency symptoms appear. Repeat application 2-4 weeks later.

FRUIT TREES AND GRAPES: Apply 1-4 pints per acre (1.25-5 liters per hectare) starting at bud break or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

VEGETABLES AND POTATOES:

Apply 1-4 pints per acre (1.25-5 liters per hectare) of **BIOMIN BORON** at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later. In greenhouses and plastic tunnels, application may be repeated everyweek or every other week.

ALMONDS, WALNUTS, PISTACHIOS AND OTHER NUT CROPS:

Apply 1-4 pints per acre (1.25-5 liters per hectare) before bloom. Repeat the application if necessary.

For best results, apply **BIOMIN BORON** when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 100 times of water.

BIOMIN BORON may be applied to fruit trees, field crops, vegetables, greenhouse crops, potatoes, grapes, citrus, and ornamental gardens.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

JH Biotech Inc.

0315-CA-B-2



MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538 , Ventura, CA 93006 U.S.A. Phone: 805-650-8933 Website: www.jhbiotech.com

Copyright © 2013 All rights reserved