ON-Gard® 5-0-0

CALIFORNIA DEPARTMENT OF FOOD AND
AGRICULTURE APR 2 0 2017

Initials: _______Date: ________

31	IAR	ΔN.	TFF	DA	IAL	YSIS
Э.		\sim 13		-	VA.	1 , 1 , 1

Total Nitrogen (N)5.0% 5.0% Water Soluble Nitrogen

Derived from: Soy Protein Hydrolysate

CDFA makes no representation as to product value, safety or use for an intended purpose! Cabeling is in accordance with the requirements of Section 14631 of the Food and Agriculture Code and Sections 2300 through 2326 of the California Code of Regulations.

ON-Gard is an innovative fertilizer based on soy protein hydrolysates. Protein hydrolysate is the organic material obtained by the hydrolysis of proteins to their constituent amino acids and short polypeptides. They are a source of nitrogen. It is rapidly absorbed by leaves and roots and translocated throughout the plant. ON-Gard is 100% water soluble and manufactured for consistency.

GENERAL INFORMATION

ON-Gard improves plant growth, supporting all phases of development

ON-Gard enhances traditional fertilizer programs to achieve high productivity. ON-Gard is non-phytotoxic and does not contain environmentally sensitive compounds. ON-Gard does not contain any animal products, animal extracts, or tannery byproducts.

USE INFORMATION AND RATES

ON-Gard is suitable for use on all growing plants. Plants respond well to ON-Gard application at key nutrient demand periods such as seed germination, bloom, fruit development, sugar assimilation, etc. ON-Gard can be applied as either a foliar spray or soil drench using any type of sprayer (hand, hydraulic, air blast, etc.) or irrigation method (overhead, overhead boom, and drip).

Field grown plants: Apply 16 – 64 fluid ounces per acre during the growth cycle of the plant. Use as much water as is needed for adequate foliage coverage or as required by any tank mix partner material(s). Applications may be repeated up to 6 times per growing season. ON-Gard can be applied as a pre-plant root dip or as a seed drench using a 1% v/v rate.

ON-Gard® Edible Field Crop Application Guide				
Crop	Foliar Application			
Field Production Fruits and Vegetables ¹	16 - 64 fl oz/ ac In sufficient water for adequate foliage coverage	2.25 - 4.5 L/ ha In sufficient water for adequate foliage coverage		

Apply up to 6 times per growing season

ON-Gard® Grape Application Guide			
	Foliar Application In sufficient water for adequate foliage coverage		
16 - 64 fl oz/ac	2.25 - 4.5 L/ha	-at fruit set	
16 - 64 fl oz/ac	2.25 - 4.5 L/ha	-at veraison	
16 - 64 fl oz/ac	2.25 - 4.5 L/ha	-tank mixed with each fungicide application	

Greenhouse, nursery, landscape, and high tunnel: On all ornamental or edible crops, apply using rates below. Use as much water as is needed for adequate foliage coverage or as required by tank mix partner material(s). ON-Gard can also be applied as a pre-plant root dip or as a seed drench using a 1% v/v rate.

ON-Gard® Application Guide					
Crop		oray Rate age to glisten)	Drench Rate (use sufficient water to penetrate root zone)		
Greenhouse/High	1 qt - 2.0 gal/100 gal	250 mL - 2.0 L/100 L	1 qt - 2.5 gal/100 gal	250 mL - 2.5 L/100 L	
Tunnel Edibles ¹	2 tsp - 2.5 fl oz/gal	2.5 mL - 20 mL/L	2 tsp - 3.3 fl oz/gal	2.5 mL - 25 ml/L	
Greenhouse,	1 qt - 2.0 gal/100 gal	250 mL - 2.0 L/100 L	1 qt - 2.5 gal/100 gal	250 mL - 2.5 L/100 L	
Nursery Ornamentals ¹	2 tsp – 2.5 fl oz/gal	2.5 mL - 20 mL/L	2 tsp - 3.3 fl oz/gal	2.5 mL - 25 ml/L	
Landscape	1 qt - 2.0 gal/100 gal	250 mL - 2.0 L/100 L	1 qt - 2.5 gal/100 gal	250 mL - 2.5 L/100 L	
Ornamentals ²	2 tsp - 2.5 fl oz/gal	2.5 mL - 20 ml/L	2 tsp - 3.3 fl oz/gal	2.5 mL - 25 ml/L	

¹ Apply every 15 days

Turf: On all types of turfgrass, apply **ON-Gard** as a drench or foliar spray. ON-Gard can be applied as a seed drench using a 1% v/v rate.

ON-Gard® Turf Application Guide				
Foliar Spra	y Rate ¹	Drench Rate ¹		
(spray on foliag	e to glisten)	(use sufficient water to penetrate root zone)		
1 qt - 2.0 gal/in up to 100 gal	2.4 – 19 L/ in up to 1000 L	1 qt - 2.5 gal/100 gal	2.4 - 23.4 L/1000L	
water/acre	water/ha	water/acre	water/ha	
1 – 6 fl oz/in up to 2 gal	24 - 190 ml/in up to 10 L	1 - 7.5 fl oz/2 gal	24 - 234 ml/10 L water	
water/1000 sq ft	water /100 sq m	water/1000 sq ft	/100 sq m	

Apply up to 6 times per growing season

Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rains are expected. Apply this product only to your lawn/garden.


SHAKE WELL BEFORE USE

ON-Gard is an excellent tank-mix partner with a wide range of products. Before tank-mixing, a jar test is recommended to evaluate physical compatibility with other products. Do not exceed recommended rate. Do not apply undiluted.

ON-Gard is stable when stored between 40.and 95 °F (4 and 35 °C). Store in a dry place. Keep out of reach of children and animals. Do not ingest. Avoid contact with eyes and skin. In case of accidental spillage of fertilizer, collect with absorbent material.

Keep out of the reach of children

NOTICE TO BUYER AND SELLER: Selter warrants that this product conforms to the description on this label and is reasonably fit for the purposes stated on this label when used and stored in accordance with the directions for use. This warranty does not extend to use of this product contrary to label directions or under conditions not reasonably foreseeable by the Selter, and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, Selter disclaims all other warranties, express or implied, including any warranty of fitness or merchantability. To the extent consistent with applicable law, Selter is abelien for consequential, special or indirect damages resulting from use or handling of this product, apply Selter's sole liability and Buyer's and User's exclusive remedy shall be limited to refund of the purchase price. This product is sold only for uses stated on its label.



Guaranteed by: BioWorks, Inc. 100 Rawson Rd, Suite 205 Victor, NY 14564 800-877-9443 www.bioworksinc.com

Container Sizes: 2.5, 5, 30 GALLONS (9.5, 19, and 113.5 LITERS)

LBS/GALLON: 9.84 @ 68° F (1.18 KG/L @ 20° C)

² Apply up to 6 times per growing season