

USE IN LOW VOLUME APPLICATORS BOTANIGARD® MAXX

GENERAL CONSIDERATIONS:

BotaniGard MAXX® can be applied effectively using low volume applicators. Increased efficacy can be obtained from these methods over that which is achieved by hydraulic spraying. Coverage is essential with BotaniGard MAXX, and when properly used through a fogger, increased contact with insect pests is possible.

TIMING & RATES

BotaniGard MAXX applications should begin at the first appearance of the insect pest (presence of eggs or small larvae/nymphs). Apply during the early morning or evening. Make 2 – 3 applications at weekly intervals. After this, switch to BotaniGard ES or 22WP formulations for applications at a longer interval (5-10 days). See rate chart on next page.

WHEN USING ULTRA LOW VOLUME APPLICATORS (FOGGERS), OBSERVE THE FOLLOWING:

- Low volume sprayers have a high degree of efficiency in coverage of leaf and stem surfaces. Thoroughly cover the plant surfaces but note that spray deposition may not be visible after application
- When using a low volume sprayer, first determine the proper application volume by spraying the material in a small area. Test the application on a few plants to assure proper coverage
- Follow equipment manufacturer instructions carefully
- Properly calibrate equipment regularly
- Ensure that nozzles are clean and not worn so that correct droplet size is applied. Smaller droplets are more desirable in a fogging situation
- Ensure that all the material is well mixed so application will be uniform throughout
- Use the maximum amount of water that a given machine allows to help ensure material is completely mixed with water
- Consider using a carrier solution appropriate to the type of fogger equipment. This will improve atomization and cause fewer large droplets to fall out in front of the fogger providing more uniform coverage. For Dramm BIO PulsFOG® equipment it is suggested to use NutriFog™ as a carrier for BotaniGard
- With most LVM equipment, keep nozzle output from directly contacting the plants. Nozzles should not be aimed directly at crop
- Unlike hydraulic or low volume mist spraying, there should be no visible evidence (leaf wetness) of the application on the plant foliage

PLANT SAFETY

Small-leaved Euphorbia varieties such as Diamond Frost®, Silver Fog, and Breathless™ may be injured with BotaniGard MAXX. Watch for these crops if foggers are used wall-to-wall in a greenhouse. When using a low volume applicator, first determine the proper application volume by applying BotaniGard MAXX in a small area. BotaniGard MAXX is an oil-based formulation that may cause injury to young plants or open blooms. Therefore, test the application on a few plants to ensure proper coverage and assess within 24 - 48 hours.

USE IN LOW VOLUME APPLICATORS BOTANIGARD® MAXX

BOTANIGARD MAXX LOW VOLUME APPLICATOR RATES: Auto-Fog™, BIO PulsFOG®, & Coldfogger™

PROGRAM	HYDRAULIC RATE/100 GAL ¹	RATE/10,000 SQ FT (DRAMM LOW 20%) ²	RATE/10,000 SQ FT (DRAMM HIGH 35%) ²	MIN. AMT. OF SOLUTION. ³ AUTO-FOG, TRACFOG, BIO PULSFOG	MIN. AMT OF SOLUTION FOR COLDFOgger
LOW LABEL RATE					
BotaniGard MAXX	32 fl oz	6.4 oz	11.2 fl oz	2 liters	10 liters
HIGH LABEL RATE					
BotaniGard MAXX	64 fl oz	12.8 fl oz	22.4 fl oz	4 liters	10 liters

1. Hydraulic sprayer rate per 100 gallons (per acre)
2. Dramm rate adjustments: Low (20%) is used for plugs and other propagation or small plants. High (35%) is used for normal finished or dense crops. These are used to bring the per acre rate down to a rate per 10,000 sq ft.
3. More solution may be used. Solution includes carrier + water. The Dramm owner's manual states that you should never use less than 2 liters of water in their foggers no matter how small an area is being fogged to ensure optimum performance of their machines. We do not recommend going below the amounts stated above. Please refer to Dramm User's Manuals for complete equipment operating instructions.
4. Maintain pH of the tank solution between pH 5.5 – 7.0

FOR MORE INFORMATION ON BOTANIGARD OR TO GET IN TOUCH WITH A REPRESENTATIVE, VISIT:

CertisBio.com



Refer to product labels for complete application details. BotaniGard® and Mycotrol® are registered trademarks of Certis Biologicals Coldfogger™, NutriFOG™, TracFOG™ and AutoFog™ are trademarks and Bio PulsFOG® is a registered trademark of the Dramm Corporation. Always read and follow label directions.

1.800.250.5024 www.CertisBio.com

